



BREAST AUGMENTATION

AFTER YOUR SURGERY

- You will likely feel drowsy upon waking up from anesthesia.
- You may have a sore throat.
- You will see/feel swelling and bruising of your breasts and flanks. You will experience a "tight" sensation in these areas.
- Usually temporary increased or loss of sensation of your nipples.
- Moderate discomfort.
- Possibly minor stinging when you urinate for the first time after surgery. This is due to the urinary catheter you will have in place during surgery.
- Sloshing sound from your breasts due to irrigation fluid and local anesthesia.
- Breasts being high and firm.
- You will have a surgery bra for support.
- There will be small pieces of tape over the incisions, eliminating the need for wound care.
- Apply ice gels to the side of your breasts for 20 minutes at a time. Do not apply heat or anything frozen directly to the skin.
- Do not sleep on your stomach.
- Rest, yet keep moving. Rest is important, although it is equally important to walk inside for 5-10 minutes every couple of hours while awake.
- Take the medication, as directed, to address any discomfort. The period of greatest discomfort does not usually last more than 24-48 hours.
- Your pain medication may constipate you. Be sure to take in plenty of good fluids such as water, broth, or sports drinks. Milk of magnesia can be purchased from the local drug store to help with constipation.
- No alcohol.
- No lifting over 10 pounds, no pushing/pulling, do not allow your heart rate above 90 bpm.
- You may remove the bandages under your arms 24 hours after surgery.
- Your postoperative visit will be scheduled for approximately 10 days after surgery.
- NO smoking. Avoid being around those who are smoking.

CALL THE OFFICE IF EXPERIENCING

- Pain not controlled by the prescribed medication.
- Excessive or sudden swelling on one side.

- Temperature higher than 101 degrees F.
- Severe nausea and vomiting.
- Pain in calf, knee, groin, or shortness of breath.
- Continuous dizziness or unusual behavior.

2 TO 7 DAYS FOLLOWING SURGERY

- During this time, you can expect to slowly progress day by day.
- Take 10 deep breaths per hour while awake. This will enhance your oxygenation and help expand your lungs.
- Please be sure to speak with Dr. Shell before you begin to drive or return to work or a regular routine at home.
- Continue a well-balanced diet with adequate hydration.
- Advice exposure to sun and heat.
- Continue to take prescribed pain medications as needed.
- Begin to transition from prescribed pain medication to over the counter alternative. Dr. Shell and staff can guide you through this as needed.
- Keep wearing provided surgery bra day and night.
- Do not sleep on your stomach.
- Avoid exercise other than walking. Leisurely daily walks will help you feel better and prevent blood clots.
- Make sure to get up and move for about 10-15 of every 60 minutes while awake.
- NO smoking. Avoid being around those who are smoking.
- At your first post-operative visit, Dr. Shell will teach you how to perform a daily breast massage that will help to maintain the surgical pocket.

1 TO 4 WEEKS FOLLOWING SURGERY

- Healing is a gradual process. It is normal to feel discouraged while waiting for swelling, bruising, discomfort, and numbness to diminish. Extra rest, a nutritious diet, and avoidance of stress are important aids to recovery. Once cleared by Dr. Shell, you may take Aspirin, Advil, Aleve, etc. at this time.
- Wash incision with Dial antibacterial soap and pat dry with clean cloth.
- The small strips of tape covering your incision will fall off 2-3 weeks after surgery.
- You may transition from your surgery to bra to a garment of your choice with no underwire.
- Once cleared by Dr. Shell, you may ease back into your workout routine.
- Continue to avoid sun exposure and heat. Use sunscreen on incision as it may tend to hyper pigment.
- NO smoking. Avoid being around those who are smoking.
- Continue with your daily breast massage.
- Do not sleep on your stomach.

4 TO 6 WEEKS FOLLOWING SURGERY

- Overall swelling continues to subside and you will begin to settle into your natural appearance. Tenderness, tightness, skin numbness, or bruising (if remaining) will continue to diminish.
- You may submerge your body in water, following complete healing of your incisions.
- You may begin wearing bras with underwire.
- You may now sleep on your stomach.
- Continue with your daily breast massage.

THE FIRST YEAR

- Continued refinements in the appearance of your breasts may take up to one full year. Your breasts will soften and settle.
- Appearances continue to change with age. Just as other areas of your body may change with age, so can your breasts.
- If you notice any signs of increasing firmness of one or both breasts, please call our office.
- Continue with your daily breast massage.
- Continue your monthly self-breast exams and mammograms as your primary care doctor recommends.
- Avoid direct sun exposure to scars the first year.
- Call our office if you notice a stitch coming up like a bristle brush along your scar. If the area around the stitch looks red and irritated, you may need antibiotics.
- Call our office if your scar becomes raised and/or lumpy.